

BIOMIN® BOOSTER V Liquid

Product Information

About BIOMIN® BOOSTER V Liquid

BIOMIN® BOOSTER V Liquid is a chelated multi-mineral which is chelated mainly by Glycine and other organic amino acids. **BIOMIN® BOOSTER V** was designed for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct trace mineral deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilisers. **BIOMIN® BOOSTER V** may be applied to all crops including fruit trees, field crops, greenhouse crops, vegetables, cotton, potatoes, onions, grapes, citrus, carrots, mangoes, banana, nuts, kiwifruit, avocado, strawberries, berries and all tropical fruit.

GUARANTEED ANALYSIS

As true amino acid Chelates: %w/w

Nitrogen (N)	1.5%
Zinc (Zn)	2.5%
Magnesium (Mg)	2.5%
Iron (Fe)	2.0%
Manganese (Mn)	2.0%
Boron (B)	0.325%
Copper (Cu)	0.65%

Magnesium, Zinc, Iron & Manganese present as Glycine chelates. Boron present as amino acid chelates. Nitrogen present as amino acid.

KEEP OUT OF REACH OF CHILDREN

PREVENTS AND CORRECTS MINERAL DEFICIENCIES

In conditions where the water and/or soil pH is alkaline, it would be advantageous to apply **MOBILIZER** to the water first before adding **BIOMIN® BOOSTER V** to the tank. To further increase availability, add **BIOMIN® BOOSTER V** to the tank when the pH of the water is 6.5 - 7.0.

Net Contents

1 Gallon 5 Gallons 1000 Litres

LOT #

ZADCO
for quality gro. pty. ltd.

Manufactured by: JH BIOTECH, INC. PO
Box 3538, Ventura, CA 93006, USA

Imported and Distributed in Australia by: **Zadco For Quality Gro PTY LTD.** ABN 57080 759 084
PO BOX 6973 Baulkham Hills BC NSW 2153 Australia T: 02 8814 6638 | F: 02 8814 6623
E: inquiries@zadco.com.au | W: www.zadco.com.au

Imported & Distributed in New Zealand by: **Roots Shoots & Fruits Ltd.** | P.O. Box 72 Waiheke Island,
Auckland New Zealand T: 09-372-9155 | M: 027-6075-764 | F: 09-372-9156 | E: rsf@rd2.co.nz

Directions and General Recommendations

Soil Application

BIOMIN® BOOSTER V may be applied before or after planting through irrigation water. The application rate is 2 - 10L / ha depending on crop, soil conditions and application methods. For maintenance, 3L / ha is recommended.

Foliar Application

ALL FIELD CROPS (grain crops, flowers, etc...): Apply 2 - 5L of **BIOMIN® BOOSTER V Liquid** per hectare when crops are 15 - 30cms high or when deficiency symptoms appear. Repeat application 2 - 4 weeks later or as needed.

ALL VEGETABLE CROPS: Apply 2 - 10L of **BIOMIN® BOOSTER V Liquid** per hectare at about 2 - 4 weeks after emergence or if deficiency symptoms appear. Repeat application 2 - 3 weeks later, or as needed.

GREENHOUSE CROPS: Apply 100 - 150ml of **BIOMIN® BOOSTER V Liquid** per 100 liters of water 2-3 weeks after emergence or transplanting, or when trace element deficiencies first appear. Repeat the application every 2-3 weeks.

ALL STONE FRUITS: Apply 2 - 10L of **BIOMIN® BOOSTER V Liquid** per hectare only as a post harvest application at least 6 weeks before dormancy.

Limited Warranty

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

APPLES, POMEFRUIT, CITRUS, AVOCADO & KIWIFRUIT: Apply 2 - 10L of **BIOMIN® BOOSTER V Liquid** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat the application 6 weeks before dormancy.

WINE & TABLE GRAPES: Apply 2 - 10L of **BIOMIN® BOOSTER V Liquid** per hectare in early spring when new leaves emerge or 2 weeks before flowering or when deficiency symptoms appear. Repeat the application 6 weeks before dormancy at a rate of 2 - 10L per hectare.

BIOMIN® BOOSTER V Liquid is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water. **BIOMIN® BOOSTER V liquid** may be used at any time throughout the season unless otherwise specified on this label. For best results and more specific recommendations, contact your nearest distributor.

Compatibility

ALWAYS run a compatibility test before application.