

# BIOMIN® BORON



## Product Information

**BIOMIN BORON** is a complexed mineral which is chelated mainly by Glycine and other organic amino acids. **BIOMIN BORON** was designed for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct BORON deficiencies and to boost crops during critical or fast growing periods.

**BIOMIN BORON** can be applied to ALL crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocadoes, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

### GUARANTEED ANALYSIS

	Dry Weight Basis %/w/w
As true amino acid Chelates:	
Boron (B)	15%

Derived as an amino acid complex.

**KEEP OUT OF REACH OF CHILDREN**

### PREVENTS AND CORRECTS MINERAL DEFICIENCIES OF BORON

In conditions where the water and/or soil pH is alkaline, it would be advantageous to apply **MOBILIZER** to the water first before adding **BIOMIN BORON** to the tank. For increased availability, add **BIOMIN BORON** to the tank when the pH of the water is 6.5 – 7.5.

**COMPATIBILITY:** Always perform a compatibility test before mixing any products with **BIOMIN BORON**.

### NET CONTENTS

2 kilograms     10 kilograms     25 kilograms

Lot #

## Directions and General Recommendations

### SOIL APPLICATION

**BIOMIN BORON** may be applied before or at planting through irrigation water. The application rate is 1.5 - 3 kilograms per hectare depending on crop, soil conditions and application methods.

### FOLIAR APPLICATION

**ALL FIELD CROPS** (grain crops, pastures, flowers, etc...): Apply 1.0 - 1.5 kilograms of **BIOMIN BORON** per hectare when crops are 15 - 30cms high or when deficiency symptoms appear. Repeat application 2 - 4 weeks later or as needed.

**ALL VEGETABLE CROPS (ESPECIALLY BRASICAS)** : Apply 1.0 - 1.5 kilograms of **BIOMIN BORON** per hectare at about 2 - 4 weeks after emergence and/or transplanting or if deficiency symptoms appear. Repeat the application 2 - 3 weeks later or as needed.

**GREENHOUSE CROPS:** Apply 100 - 150 grams of **BIOMIN BORON** per 100 liters of water 2-3 weeks after emergence or transplanting, or when trace element deficiencies first appear. Repeat the application as needed.

**APPLES:** Apply 0.5 - 2.0 kilograms of **BIOMIN BORON** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat the application post harvest at 6 weeks before dormancy at a rate of 0.5 - 2.0 kilograms per hectare.

### LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

**FRUIT TREES & AVOCADOS:** Apply 0.5 - 2.0 kilograms of **BIOMIN BORON** per hectare in early spring when new leaves emerge and / or 2 weeks before flowering. Repeat the application at any time throughout the season and post harvest at 6 weeks before dormancy at a rate of 0.5 - 2.0 kilograms per hectare.

**KIWIFRUIT:** Apply 0.5 - 2.0 kilograms of **BIOMIN BORON** per hectare in early spring when new leaves emerge or 2 weeks before flowering. Repeat the application post harvest at 6 weeks before dormancy at a rate of 0.5 - 2.0 kilograms per hectare.

**GRAPES:** Apply 0.5 - 2.0 kilograms of **BIOMIN BORON** per hectare in early spring when new leaves emerge or 2 weeks before flowering and / or when deficiency symptoms appear. Research has shown that an application of Boron at veraison can aid with sugar accumulation. Repeat the application of **BIOMIN BORON** post harvest at 6 weeks before dormancy at a rate of 0.5 - 2.0 kilograms per hectare.

**BIOMIN BORON** is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water. **BIOMIN BORON may be used at any time throughout the season.** For best results and more specific recommendations, contact your nearest distributor.

Imported and Distributed in Australia exclusively by: Zadco For Quality Gro PTY LTD. ABN 57080 759 084 | 9/9A Foundry Road, Seven Hills NSW 2147  
Ph: 02 9838 9111 Fax: 02 9838 9110 | email: inquiries@zadco.com.au. website: www.zadco.com.au Imported and Distributed in New Zealand exclusively by:  
Roots Shoots and Fruits Ltd | PO Box 72, Waiheke Island, Auckland, New Zealand | Ph: +64(0) 9372 9155 Fax: +64(0) 9372 9156 | email: rsf@rd2.co.nz  
website: www.rd2.co.nz Manufactured by: JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA